

## Evaluation of antifungal, antioxidant and phytochemical screening of *Ziziphus jujuba* Miller leaves from the northwestern region of Algeria

Zoubir Belmokhtar\*<sup>1,4</sup>, Merad Yassine<sup>2</sup>, Harrachi Abdelkader<sup>3</sup>, Farsi Zohra<sup>4</sup>, Ghenim Samira<sup>4</sup>, Mermouri Lamia<sup>1</sup>, Mezmez Ryad<sup>1</sup> and Kaid Harche Meriem<sup>1</sup>

<sup>1</sup>\*Laboratory of Plant and Microbial Productions and Valorization (LP2VM), Department of Biotechnology, University of Science and Technology of Oran: Mohamed Boudiaf, El Mnaouar, BP 1505, Bir El Djir 31000, Oran, Algeria, <sup>2</sup>Department of Medicine, Faculty of Medical Sciences, University Djillali Liabes, BP 89 Sidi Bel Abbes 22000, Algeria, <sup>3</sup>Laboratory of Microbiology, University Hospital Center of Sidi Bel Abbes BP 89 Sidi Bel Abbes 22000, Algeria and <sup>4</sup>Department of Environmental Sciences, Faculty of Natural Science and Life, University Djillali Liabes, BP 89 Sidi Bel Abbes 22000, Algeria

### Abstract

*Ziziphus jujuba* is used in Algeria in various herbal remedies. Unfortunately this species is little known phytochemically. In order to valorize medicinal plants grown in northwestern Algeria, we are interested to evaluating the antioxidative and antifungal activities of *Ziziphus jujuba*, and to determine the main phytochemical classes involved in these activities, to improve the therapeutic knowledge of this species. Phytochemical screening was performed using standard methods. Quantification of total polyphenols, flavonoids and condensed tannins was performed by colorimetric techniques using a UV-visible spectrophotometer. The extract was further evaluated for their antifungal activity against seven strains of pathogenic fungi, by well microdilution method, and the free radical scavenging activity of extract was measured by using the DPPH. The results of phytochemical screenings showed the presence of flavonoids, alkaloids, saponins, gallic tannins, leucoanthocyanins and cardiac glycosides. Quantitative analysis of phenolic compounds shows that total polyphenols, flavonoids and condensed tannins have 240.78 (mg GAE/g DE); 113.6 (mg EC/g DE) and 78.02 (mg EC/g DE) respectively. The evaluation of the antioxidative activity, show that the extract of *Ziziphus jujuba* leaf has an IC<sub>50</sub> of 0.38 mg/ml and at a dose of 1 mg/ml reduces 91.78% of oxidized DPPH in a very short time. Among the seven fungal pathogenic strains used for the test of antifungal activity, the most potent activity was against *Candida albicans* with a MIC of 0.0975 mg/ml. *Ziziphus jujuba* leaves contain different biologically active molecules which justify their use in phytotherapy.

**Key words:** *Ziziphus jujube*, Phytochemical, Antifungal, DPPH

Full length article \*Corresponding Author, e-mail: [zoubir\\_31@yahoo.fr](mailto:zoubir_31@yahoo.fr)

### 1. Introduction

*Ziziphus jujuba* locally called Sfisef in northwestern Algeria, belongs to the family Rhamnaceae, and is a thorny tree 4 to 10 m long. The leaves are 4 to 8 cm long, ovate to elliptic in shape with finely toothed edges, bright green in color, but yellow in autumn before falling. Small yellow flowers develop in the axils of the leaves in early summer. The fruit is an edible drupe, round to elongated, smooth and green before maturity, it folds and becomes mahogany or reddish-brown at the final stage of maturity, carries a single seed.

In traditional medicine, Jujube is introduced as a blood purifier, sedative, expectorant, cough suppressant, anti-asthmatic, laxative, wound healer, anaphrodisiac. It also contributes to the treatment of rectal and intestinal

ulcers/diseases as well as liver diseases. Ripe jujube fruit has laxative properties, but unripe jujube heals diarrhea [1]. Their fruits and seeds are usually applied in traditional Chinese medicine for the treatment of various diseases, such as anorexia, lassitude, insomnia, anxiety [2]. The leaves have been also used as a folk medicine to treat children suffering from typhoid fever, furuncle and eczema in China [3].

Currently, several laboratory work that can be in vivo or in vitro confirms that this species has interesting biological activities such as anti-inflammatory [4], anti-diarrhoeal [5], sedative [6], anticancer [7], antibacterial [8], antiobesity [9], antidiabetic [10] and hepatic protective [11-13].

In Algeria *Zizyphus jujuba* is very little known chemically, so no work is done on the antifungal activity on this species. As part of the valuation of medicinal species, we are interested in this study to identify molecules of secondary metabolism by chemical screening, quantify polyphenols, flavonoids and tannins, and evaluate the antioxidative and antifungal activity of this species which develops in northwestern Algeria.

## 2. Materials and Methods

### 2.1 Plant Material

The leaves of *Zizyphus jujuba* are collected from the region of Sfisef, this region takes the vernacular name of the species studied because of their abundance. It is located in the province of Sidi Bel Abbes in northwestern Algeria. The species has been identified by the forest protection directorate of the region. The leaves were dried out of light and moisture at room temperature and then reduced to a fine powder.

### 2.2 Phytochemical Screening

Phytochemical analysis of the extracts was performed according to standard methods with some modifications [14-16]. Leaves of *Zizyphus jujuba* are screened for the following phytoconstituents: alkaloids, tannin, saponins, terpenoids, flavonoids, cardiac glycosides, quinones, anthraquinones, anthocyanins and leucoanthocyanins.

#### 2.2.1 Test for Flavonoids (Cyanidin Reaction)

To 5 ml of infusion, were added 5 ml of hydrochloric alcohol, 0.5 g of magnesium chips and 1 ml of isoamylic alcohol which collect the red, orange or violet color when there is presence of flavonoids.

#### 2.2.2 Test for Leucoanthocyanins

The same reaction of cyanidin cited above but without addition of magnesium turnings. After 15 min of heating in a water bath, the appearance of red, cherry-red, brown or purplish red color indicates the presence of leucoanthocyanin.

#### 2.2.3 Test for Alkaloids

25 mg of extract was dissolved in 5 ml of 5% aqueous HCl and filtered. These filtrates were divided into three equal parts. Drops of Wagner, Mayer and Dragendorff reagents were added to each. A red-brown precipitate (Wagner), yellowish-white precipitate (Mayer) and red-orange precipitate (Dragendorff) indicated the presence of alkaloids.

#### 2.2.4 Saponins (Foam Test)

1 ml solution of extract was diluted with distilled water to 20 ml and shaken vigorously and set aside for 15 min. Development of stable foam suggests the presence of saponins.

##### 2.2.4.1 Determination of Foaming Index

For determination of foaming index, 1 g of aerial parts was taken in a 250 ml conical flask containing 100 ml of boiling water. The moderate boiling was maintained for

30 min. It was cooled and filtered into a 100 ml volumetric flask and the volume was readjusted to 100 ml with distilled water. In 10 stoppered test tubes (height 16 cm, diameter 16 mm), the solution was taken in successive portion of 1, 2, 3, up to 10 ml. The volume of the solution in each test tube was adjusted with distilled water to 10 ml. The tubes were corked and shaken in a lengthwise motion for 15 sec, two shakes per second was maintained in each tube. The test tubes were allowed to stand for 15 min and height of the foam was measured. Foaming index =  $1000 \frac{a}{v}$ ; Where, a= the volume in ml of the decoction used for preparing dilution in the tube where foaming to a height of 1 cm is observed.

#### 2.2.5. Test for Tannins

2–3 drops of FeCl<sub>3</sub> 1% solution was added to 5 ml of infusion in water. A blackish-blue color indicated the presence of Gallic tannins, while a greenish-black color indicated the presence of catechol tannins.

#### 2.2.6. Test for Cardiac Glycoside (Keller-Killiani Test)

5 ml of each extract was treated with 2 ml of glacial acetic acid containing a few drops of ferric chloride and 1 ml of H<sub>2</sub>SO<sub>4</sub> along the side of the test tube. The formation of brown ring at the interface gives positive indication for cardiac glycoside and a violet ring may appear below the brown ring.

#### 2.2.7. Test for Quinones

Concentrated sulphuric acid (1 ml) was added to 1 ml of plant extract. The formation of red color indicates the presence of quinones.

#### 2.2.8. Test for Anthraquinone (Borntrager Test)

Powdered drug is mixed with chloroform or any water immiscible solvent which is filtered and to the filtrate, adds ammonia. The ammoniacal layer shows after shaking pink red or violet color due to presences of anthraquinone.

#### 2.2.9. Test for Anthocyanins

Anthocyanins were revealed by adding 5 ml of 5% infused with 5 ml of H<sub>2</sub>SO<sub>4</sub> 10% and 5 ml of NH<sub>4</sub>OH 50%. The color of the infused is accentuated by acidification, then turns blue in basic medium in the presence of anthocyanins.

## 2.3 Preparation of the Hydroethanolic Extract

10 g of powder were exhaustively three times extracted with ethanol 70% at a ratio of 1:10 (w/v) using the maceration method for 12 h at ambient temperature. The extracts were collected and concentrated using a rotary vacuum evaporator at 40°C then the solid crude extract was kept in the refrigerator (4°C) for further analysis.

## 2.4 Quantification of Phenolic Compounds

### 2.4.1 Determination of Total Polyphenol Content

The total phenolic contents are determined by the Folin-Ciocalteu method [17]. 20 µl of extract (1 mg/ml) was mixed with 1.58 ml water, and then added 100 µl of the Folin-Ciocalteu (1/10) and 300 µl of Na<sub>2</sub>CO<sub>3</sub> (7.5%). After 90 min at room temperature, the absorbance was measured at 765 nm. The total polyphenol content was determined from extrapolation of calibration curve which was made by

preparing gallic acid solution (10 to 100 mg/ml). The estimation of the phenolic compounds was carried out in triplicate. The total phenolic contents were expressed as milligrams of gallic acid equivalents (GAE) per g of dried extract (DE).

#### 2.4.2 Determination of Total Flavonoids Content

The total flavonoids content of leaves extract was estimated by the method that uses aluminium chloride [18]. 1 ml of sample containing 1 mg/ml of dry extract was mixed with 5 ml of distilled water and subsequently with 0.3 ml of a NaNO<sub>2</sub> solution (5%) (Merck). After 6 min, 0.3 ml of 10% AlCl<sub>3</sub>.6H<sub>2</sub>O (Merck) was added and allowed to stand for 6 min, then 2 ml of 1 mol/l NaOH was added to the mixture. Immediately, water was added to bring the final volume to 10 ml and the mixture was thoroughly mixed and allowed to stand for another 15 min. The absorbance of the mixture was then determined at 510 nm. Flavonoid contents were calculated using a standard calibration curve developed by using the known concentrations of catechin (10–100 mg/ml). The results were expressed as mg of catechin equivalent (EC) per g of dried extract (DE).

#### 2.4.3 Determination of Condensed Tannin Content

The condensed tannin contents were determined by a modification of the vanillin/HCl method [19]. 100 µl of extract (1 mg/ml) was pipetted into a test tube and 5 ml vanillin/HCl reagent was added. The reaction was carried out in the dark at room temperature for 20 min, and then absorbance was measured at 500 nm.

Condensed tannin contents were calculated using a standard calibration curve developed by using the known concentrations of catechin (10–100 mg/ml). The results were expressed as mg of catechin equivalent (EC) per g of dried extract (DE).

### 2.5 Study of Antioxidant Activity

#### 2.5.1 Scavenging Activity Against 1,1-Diphenyl-2-Picryl Hydrazyl Radical

The antioxidant activity of the samples and standards was determined by way of the radical scavenging activity method using 2,2-diphenyl-1-picrylhydrazyl radical (DPPH). 0.1 ml of the sample at different concentrations (0.05–10 mg/ml) was each added to 3.9 ml of DPPH methanolic solution (60 µmol/l). The blank sample consisted of 0.1 ml of methanol added to 3.9 ml of DPPH. After a 30 min of incubation period at room temperature in the dark, the absorbance was measured at 517 nm [20].

$$\% \text{ Scavenging Activity} = [(A_0 - A_1)/A_0] \times 100$$

Where, A<sub>0</sub> is the absorbance of the control and A<sub>1</sub> is the absorbance of the extract. IC<sub>50</sub> value and the percent inhibition of DPPH are calculated from the graph. Ascorbic acid was used for comparison.

#### 2.5.2 Kinetic Behaviour of the DPPH radical-scavenging activity

In this study, we examined the kinetics of the chemical reaction between the DPPH and the extract at a

concentration of 1 mg/ml every 5 min for 30 min, compared to a blank by replacing the extract with methanol. Ascorbic acid at a concentration of 1 mg/ml is used for comparison.

### 2.6 Antifungal Activity

#### 2.6.1 Fungal Strain

Seven pathogenic strains of fungal isolates were provided by the laboratory of mycology and parasitology from the university hospital center of Sidi Bel Abbas (Algeria), all isolates were identified using growth and colony characteristics, morphology, germ tube formation and the sugar fermentation, according to standard methods. They were maintained in culture on Sabouraud glucose (4%) agar medium. Fungal isolates and their origins are grouped in Table 1.

#### 2.6.2 Minimum inhibitory concentration against different fungal strains

The minimum inhibitory concentration (MIC) was determined in sterile 96-well microplates using the broth microdilution method, by the technique described by [21] with some modifications.

Aliquot of the crude hydroethanolic leaves extract was dissolved in DMSO 10% at concentrations of 100 mg/mL and sterilized by filtration.

Fungal culture was suspended in 5mL sterile 0.85% NaCl. Turbidity was adjusted to Mc Farland 0.5 scale, equivalent to 1.5×10<sup>6</sup> (CFU/mL).

Each well contained 50 µL of serial double dilutions of the extract (in a Sabouraud broth medium with concentrations that range from 100 mg/ml to 0.0487 mg/ml) is inoculated aseptically with 50 µl of inoculum suspension. The control well containing broth Sabouraud medium without extract was inoculated only with fungal suspension. The plates were incubated at 28°C for 3 days and read visually.

The MIC: It is the lowest concentration of the sample at which the tested microorganisms did not demonstrate any visible growth after incubation.

### 2.7 Statistical Analysis

All determinations were made in the triplicate and the values were averaged and reported along with the standard deviation (±). Statistical analysis was carried out by using Microsoft Excel 2010 software.

## 3. Results and Discussion

### 3.1 Phytochemical Screening

The results of the phytochemical screening of *Zizyphus jujube* are summarized in Table 2.

Phytochemical tests carried out on *Zizyphus jujuba* leaves show the presence of flavonoids, alkaloids, saponins, gallic tannins, leucoanthocyanins and cardiac glycosides, and absence of quinones, anthraquinones and anthocyanins.

The phytochemicals compounds detected are known to be of medicinal importance, can protect and lower the risk against several chronic diseases [22-23].

We found that the saponin foam index is high, which means that the leaves of this species are a very important source of these molecules. Saponins have been found to possess significant anti-cancer, antimicrobial, hypocholesterolemic, hypoglycemic, immuno-stimulant and anti-inflammatory properties [24-28].

This species is known as a sedative, this activity results from the hypnotic effect of saponins such as jujubosides which may be a good source of main compounds for new hypnotics, and the serotonergic system may involve in the hypnotic effect of jujubosides [29].

Cardiac glycosides are used for congestive heart failure and cardiac arrhythmias treatment; however, *Zizyphus jujuba* has been used in phytotherapy to treat asthma [1].

Flavonoids, phenolic acids and caffeic acid were isolated and identified by NMR and mass spectrometry in *Zizyphus oxyphylla*; a species of the same genus of jujuba [30].

### 3.2 Total Phenolic, Flavonoids and Tannin Contents

The results of quantitative analysis are represented as a histogram in Figure 1 and summarized in the table 3.

The total phenolic, flavonoids and tannin content of hydro-ethanolic extracts are 240.78 (mg GAE/g DE); 113.6 (mg EC/g DE) and 78.02 (mg EC/g DE) respectively.

In comparison with other work on the same specie [31-32] we noticed that they found different amounts of these molecules.

This difference results from several factors: the geographical position, the powder/solvent ration, the extraction technique, the method used for the quantification and the standard used.

Other works show that the flavonoids were believed to be the major bioactive components in jujube leaves, which have been shown to be responsible for cardio-protective, anticancer, antidiabetic, anti-aging and neuro-protective effects [33-34]. And the best sampling time is June so that it is a better new resource for health promotion based on flavonoids of *Zizyphus jujuba* leaves [3].

### 3.3 Study of Antioxidant Activity

The DPPH radical scavenging ability of the hydro-ethanolic extract of *Zizyphus jujuba* leaves in comparison with that of ascorbic acid are shown in Table 3.

The ability to trap DPPH with the hydro-ethanolic extract of *Zizyphus jujuba* leaf was slightly lower than that of ascorbic acid at doses ranging from 0.1 to 1 mg / mL. Therefore, IC<sub>50</sub> of *Zizyphus jujuba* leaf extract is slightly higher (0.38 mg/ml) than that of ascorbic acid (0.31 mg/ml). At a dose of 1 mg/ml, *Zizyphus jujuba* leaf extract reduces 91.78% of oxidized DPPH (Figure 2).

Other studies show that, the DPPH inhibition capacity varies between 90 and 98% of raw leaf extracts of *Zizyphus jujuba* from Oman, obtained from different solvents [35], and it is between 70.69% and 93.93% in fruit

extract from different regions of Iran [32], and it is between 33.6 and 98.6 in different Chinese jujube obtained from different station [36].

The kinetics study of the hydroethanolic extract of the leaves of *Zizyphus jujuba* at a concentration of 1 mg/ml against the DPPH, shows that the reaction is fast and almost instantaneous. The color change that expresses the passage of DPPH from the radical form (DPPH•) at the stable non-radical reduced form (DPPH-H) is done in the first five minutes, then the equilibrium state is reached immediately and the reduction is almost complete (Figure 3).

*Zizyphus jujuba* leaf infusion contains rich sedative flavonoids and antioxidants. These findings suggest that this infusion has good radical scavenging ability by reducing the risk of oxidative damages and it can very healthy beverage for the public [33]. Moreover, [37] found that *Zizyphus jujuba* contains antioxidant molecules that protect the body from oxidative stress induced by H<sub>2</sub>O<sub>2</sub> and also protect it from neurodegeneration. A recent studies shows that eight flavonoids were identified and quantified as kaempferol and quercetin glycosides, which represent a stronger DPPH and ABTS radical-scavenging activities in a much shorter time [34] and anti-proliferative activities [38].

### 3.4 Antifungal Activity

The result of the antifungal activity is shown in figure 4, and the minimum concentrations of inhibition on each fungal species are grouped in the table 4.

The hydro-ethanolic leaves extract shows antifungal activity against all fungi tested but with different concentration.

The leaf extract of *Zizyphus jujuba* showed antifungal activity against yeast *Candida albicans* and filamentous fungi such as *Aspergillus* sp and *Rhizopus* sp., and dermatophytes like *Trycophyton* sp. and *Microsporium*.

The maximum activities have been found against *Candidas* (0.0975 mg/ml), *Aspergillus niger* (0.39 mg/ml) and *Microsporium canis* (3.12 mg/ml).

Both *Alternaria* sp. and *Penicillium* sp. have been eliminated with the same concentration of the extract 12.5 mg/ml.

However the lowest activity is determined against *Rhizopus* sp. with minimum inhibitory concentration of 50 mg/ml.

In comparison with other studies on the same specie, the ethanolic extract of *Zizyphus jujuba* fruit showed a MIC of 2.35 mg/ml against *Candida albicans* and 2.86 mg/ml against *Aspergillus fumigatus* [39]. Further work indicates that the methanolic extract of *Zizyphus jujuba* leaves has MICs of 15 mg/ml against *Aspergillus flavus*, 20 mg/ml against *Tricophyton rubrum* and 35 mg/ml against *Candida albicans* [40].

The phytochemical analysis of the leaf extracts showed the presence of saponnins, tanins, flavonoids and

alkaloids, probably this antifungal activity resulting of those phytocompounds.

Phytoconstituents employed by plants to protect them against pathogenic microorganisms. The antimicrobial potency of the plant is attributed to the chemical structure and concentration of their active constituents.

Many polyphenols, such as simple phenols, flavonoids and tannins have antifungal activity, which depends in part on the number and position of the hydroxyl groups. They can inhibit the enzymes of microorganisms by reacting with sulfhydryl groups or form complexes with extracellular and soluble proteins; they could also disrupt cell membranes [41-42]. Another studies showed that tannins are toxic to filamentous fungi and yeasts showed that low tannin concentrations altered the morphology of *Crinipellis pernicioso* germ tubes [43-44]. In contrast, a

previous study in Pakistan indicated that *Zizyphus jujuba* lack potent antifungal activity, no effect was recorded against *Aspergillus flavus* while a moderate antifungal activity was described against *Aspergillus niger* [45].

Saponins are stored in plant cells as inactive precursors, but are readily converted into biologically active antibiotics by enzymes in response to pathogen attack. They appear to act by disrupting the membrane integrity of fungal cells [41].

The alkaloids can exhibit antimicrobial activity by many mechanisms, DNA intercalation, targeting RNA polymerase, gyrase and topoisomerase IV, inhibition of cell division [46-48], perturbations in the biosynthesis or metabolism of heme [49], and inhibiting enzyme activity [50].

Table 1: List of fungal strains used and their origins

Clinical fungal strains	Source
<i>Candidas</i>	Oral mycosis
<i>Aspergillus niger</i>	Ear discharge
<i>Rhizopus sp</i>	Hospital air sample
<i>Alternaria sp</i>	Hospital air sample
<i>Penicillium sp</i>	Tinea unguium
<i>Trichophyton rubrum</i>	Tinea corporis
<i>Microsporum canis</i>	Tinea capitis

Table.2 Qualitative phytochemical analysis of *Zizyphus jujba* leaves

Flavonoïds	+
Alkaloids	+
Saponins	+
Foam Index	1000
Tannin (FeCl <sub>3</sub> )	+ Gallic
Leucoanthocyanins	+
Cardiac Glycosides	+
Quinones	-
Anthraquinones	-
Anthocyanins	-

Table 3: IC50 values and the percentage inhibition of DPPH of hydro-ethanolic leaf extract of *Zizyphus jujuba* and ascorbic acid

Sample	IC <sub>50</sub> (mg/ml)	% DPPH radical scavenging activity at 1mg/ml
Hydro-ethanolic extract of <i>Zizyphus jujba</i> leaves	0.38	91, 78
Ascorbic acid	0.31	99.98

Table 4: Minimum inhibitory concentrations values of hydro-ethanolic leaves extract of *Zizyphus jujuba* against clinical fungal isolates

Clinical Fungal Strains	MIC (mg/ml)
<i>Candidas</i>	0.0975
<i>Aspergillus niger</i>	0.39
<i>Rhizopus sp</i>	50
<i>Alternaria sp</i>	12.5
<i>Penicillium sp</i>	12.5
<i>Trichophyton rubrum</i>	6.25
<i>Microsporum canis</i>	3.12




Figure 1: Total phenolic, flavonoid, and condensed tannin contents of hydro-ethanolic extract of *Zizyphus jujuba* leaves. Total phenolic were expressed by mg of GAE/g of DE. Total flavonoid and condensed tannin contents were expressed by mg of CE/g of De




Figure 2: Radical scavenging activity for leaves extracts of *Zizyphus jujuba* and ascorbic acid at different concentrations. The figure shows the percentage inhibition of DPPH and the IC50.




Figure 3: Kinetic behavior of antioxidant activity of hydro-ethanolic extract of *Zizyphus jujuba* leaves against DPPH, compared with that of ascorbic acid and blank




Figure 4: Micro-dilution plates that represent the minimum inhibitory concentrations (MIC) of hydro-ethanolic leaves extract of *Zizyphus jujuba* by Broth microdilution method, against different isolated pathogenic fungal strains

#### 4. Conclusion

It can be concluded that *Zizyphus jujuba* contains various important bioactive compounds belonging to the different phytochemical classes, which justifies their use as a medicinal plant. This species can be used against pathogenic fungi and oxidative damage. Further studies are needed to isolate active biomolecules from leaf extract and elucidate its mechanism of action.

#### References

- [1] A. Ghobadi, F. Amini-Behbahani, A. Yousefi, M. Taghavi Shirazi and N. Behnoud. (2019). Medicinal and Nutritional Properties of *Zizyphus jujuba* Mill. in Traditional Persian Medicine and Modern Phytotherapy . Crescent Journal of Medical and Biological Sciences 6 (2): 146–150.
- [2] S. Guo, J.A. Duan, Y. Tang, Y. Qian, J. Zhao, D. Qian, S. Su, E. Shang. (2011). Simultaneous qualitative and quantitative analysis of triterpenic acids, saponins and flavonoids in the leaves of two *Zizyphus* species by HPLC-PDA-MS/ELSD. Journal of Pharmaceutical and Biomedical Analysis 56: 264– 270.
- [3] X. Cui, Z. Ma, L. Bai, Y. Wu, S. GUO , Q. Liu, L. Zhang, C.T. Ho and N. Bai. (2017). Phytochemical analysis of *Zizyphus jujuba* leaves in six cultivars at the whole life stage by high performance liquid chromatography. Chemical Research in Chinese Universities 33 (5): 702–708.
- [4] S. Kumar, M.S. Ganachari, Banappa and V.S. Nagoor. (2004). Anti-inflammatory activity of *Zizyphus jujuba* Lam leaves extract in rats. Journal of natural remedies 4 (2): 183 – 185.
- [5] A.M. Mesaik, H.W. Poh, O.Y. Bin, I. Elawad and B. Alsayed. (2018). In Vivo Anti-Inflammatory, Anti-Bacterial and Anti-Diarrhoeal Activity of *Zizyphus Jujuba* Fruit Extract. Open Access Macedonian Journal of Medical Sciences 6 (5): 757-766.
- [6] J.G. Jiang, X.J. Huang, J. Chen and Q.S. Lin. (2007). Comparison of the sedative and hypnotic effects of flavonoids, saponins, and polysaccharides extracted from Semen *Zizyphus jujube*. Natural Product Research 21 (4): 310-320.
- [7] S. Ebrahimi, H. Mollaei and R. Hoshyar. (2017). *Zizyphus Jujube*: a review study of its anticancer effects in various tumor models invitro and in vivo. Cellular and Molecular Biology 63 (10): 122-127.
- [8] S.H. Abd-Alrahman, M.M.S. Bekhit, M.E.A. Elhalwagy, W.M. Abdel-Mageed and A.A. Radwan. (2013). Phytochemical Screening and Antimicrobial Activity of EthOH/Water *Zizyphus jujuba* Seeds Extracts. Journal of pure and applied microbiology 7 (Spl. Edn.): 823-828.
- [9] H. Kubota, R. Morii, A. Kojima-Yuasa, X. Huang, Y. Yano, I. Matsui-Yuasa. (2009). Effect of *Zizyphus jujuba* extract on the inhibition of adipogenesis in 3T3-L1 preadipocytes. The American Journal of Chinese Medicine 37 (03): 597-608.
- [10] Z. Sherdil, H. Madni, R. Mirbadalzadeh. (2009). Investigation into the hypoglycemic effect of hydroalcoholic extract of *Zizyphus Jujuba* Leaves on blood glucose and lipids in Alloxan-Induced diabetes in rats. Iranian Journal of Diabetes and Lipid Disorders 8 (1): 13-19.
- [11] P.K. Mukherjee, K. Maiti, K. Mukherjee and P.J. Houghton. (2006). Leads from Indian medicinal plants with hypoglycemic potentials. Journal of Ethnopharmacology 106: 1-28.
- [12] C. F. Chen, J.F. Lee, D. Wang, C.Y. Shen, K.L. Shen and M.H. Lin. 2010. Water Extract of *Zizyphus Jujube* Attenuates Ischemia/Reperfusion-Induced Liver Injury in Rats (PP106). Transplantation Proceedings 42: 741-743.
- [13] A.K. Waqar, M. Naveed, K. Haroon and A. Rauf .(2014). Pharmacological and Phytochemical

- Studies of Genus *Zizyphus*. Middle-East Journal of Scientific Research 21 (8): 1243-1263,
- [14] J.B. Harborne. (1973). *Phytochemical Methods*. London, UK: Chapman and Hall Ltd.
- [15] G.E. Trease and W.C. Evans. (1989). *Pharmacognosy*. 13th Edn. London: Baillière Tindall.
- [16] Z. Belmokhtar, Y. Merad, A. Chaib, W. Ouchen, R. Mezmez and M. Kaid-Harch. *Phytochemical Screening and the Effect of Temperature Treatment on the Extraction Yield of Phenolic Compounds and the Antioxidative Activities of Two Medicinal Species Growing in Algeria: *Ammi visnaga* and *Mentha pulegium**. *Research & Reviews: A Journal of Pharmacognosy* 5(3):18-25.
- [17] V.L. Singleton and J.A. Rossi. 1965. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *American Journal of Enology and Viticulture* 16 (3): 144–158.
- [18] V. Dewanto, X. Wu, K.K. Adom and R.H. Liu. (2002). Thermal Processing Enhances the Nutritional Value of Tomatoes by Increasing Total Antioxidant Activity. *Journal of Agricultural and Food Chemistry* 50 (10): 3010–3014.
- [19] R.E. Burns. (1971). Method of Estimation of Tannin in Grain Sorghum. *Agronomy Journal* 63 (3): 511-512.
- [20] W. Brand-Williams, M.E. Cuvelie and C. Berset. (1995). Use of a Free Radical Method to Evaluate Antioxidant Activity. *LWT Food Science Technology* 28 (1): 25–30.
- [21] M. Balouiri, M. Sadiki and S.K. Ibensouda. (2016). Methods for in vitro evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis* 6 (2): 71-79.
- [22] N. El-Najjar, H.A. Gali-Muhtasib, R. Ketola, P. Vuorela, A. Urtti and H. Vuorela. (2011). The Chemical and Biological Activities of Quinones: Overview and Implications in Analytical Detection. *Phytochemistry Reviews* 10: 353–370.
- [23] S. Ogawa and Y. Yazaki. (2018). Tannins from *Acacia mearnsii* De Wild. Bark: Tannin Determination and Biological Activities. *Molecules* 23 (4): 837.
- [24] P. Navarro, R.M. Giner, M.C. Recio, S. Manez, M. Cerda- Nicolas and J.L. Rios. (2001). In vivo anti-inflammatory activity of saponins from *Bupleurum rotundifolium*, *Life Sciences*. 68 (10): 1199–1206.
- [25] S. Man, W. Gao, Y. Zhang, L. Huang and C. Liu. (2010). Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia* 7: 703-714.
- [26] G. Francis, Z. Kerem, H.P. Makkar, K. Becker. (2002). The biological action of saponins in animal systems: A review. *British Journal of Nutrition* 88: 587–605.
- [27] M. Marrelli, F. Conforti, F. Araniti and G.A. Statti. (2016). Effects of saponins on lipid metabolism: A review of potential health benefits in the treatment of obesity. *Molecules* 21 (10): 1404.
- [28] Z. Jin, L. Gao, L. Zhang, T. Liu, F. Yu, Z. Zhang, Q. Guo and B. Wang. (2017). Antimicrobial activity of saponins produced by two novel endophytic fungi from *Panax notoginseng*. *Natural Product Research* 31 (22): 2700-2703.
- [29] J.X. Cao, Q.Y. Zhang, S.Y. Cui, X.Y. Cui, J. Zhang, Y.H. Zhang, Y.J. Bai and Y.Y. Zhao. (2010). Hypnotic effect of jujubosides from Semen *Ziziphi Spinosae*. *Journal of Ethnopharmacology*, 130 (1): 163-166.
- [30] M. Zubair, K. Rizwan, N. Rasool and R.B. Tareen. 2018. Isolation of secondary metabolites from *Zizyphus oxyphylla* Fruit. *International Journal of Chemical and Biochemical Sciences*. 13: 27-30.
- [31] M. Elalouia, A. Laamouria, A. Ennajaha, M. Cerny, C. Mathieub, G. Vilaremb, H. Chaarc and B. Hasnaoui. (2016). Phytoconstituents of leaf extracts of *Zizyphus jujuba* Mill. plants harvested in Tunisia. *Industrial Crops and Products* 83: 133–139.
- [32] A. Azizi and M. Pirbodaghi. (2016). Regional variations of antioxidant capacity and phenolic properties in the Iranian jujube collection. *Journal of Herbal Drugs* 6 (4): 199-209.
- [33] R. Zhang, J. Chen, Q. Shi, Z. Li, Z. Peng, L. Zheng and X. Wang. (2014). Phytochemical analysis of chinese commercial *Zizyphus*, jujube leaf tea using high performance liquid chromatography–electrospray ionization-time of flight mass spectrometry. *Food research international* 56: 47–54.
- [34] L. Zhang, P. Liu, L. Li, Y. Huang, Y. Pu, X. Hou and L. Song. (2018). Identification and antioxidant activity of flavonoids extracted from Xinjiang Jujube (*Zizyphus jujuba* Mill.) leaves with ultra-high pressure extraction technology. *Molecules* 24 (1): 122.
- [35] A. Al-Saeedi, M.T.H. Al-Ghafri and M.A. Hossain. (2016). Comparative evaluation of total phenols, flavonoids content and antioxidant potential of leaf and fruit extracts of Omani *Zizyphus jujuba* L. *Pacific Science Review A: Natural Science and Engineering* 18 (1): 78–83.
- [36] J.W. Li, S.D. Ding, X.L. Ding. (2005). Comparison of antioxidant capacities of extracts from five cultivars of Chinese jujube. *Process Biochemistry* 40: 3607-3613.

- [37] A. Küçükgül. 2016. The Antioxidant Effects of *Ziziphus Jujuba* on Neurodegeneration. *Etilik Veteriner Mikrobiyoloji Dergisi*, 27 (2), 108-112.
- [38] L. Song, P. Liu, Y. Yan, Y. Huang, B. Bai, X. Hou and L. Zhang. (2019). Supercritical CO<sub>2</sub> fluid extraction of flavonoid compounds from Xinjiang jujube (*Ziziphus jujuba* Mill.) leaves and associated biological activities and flavonoid compositions. *Industrial Crops & Products* 139: 111508.
- [39] F. Daneshmand, H. Zare-Zardini, B. Tolueinia, Z. Hasani and T. Ghanbari. (2013). Crude Extract from *Ziziphus Jujuba* Fruits, a Weapon against Pediatric Infectious Disease. *Iranian Journal of Pediatric Hematology Oncology* 3 (1): 216-221.
- [40] A.L. Abubakar, A. Dandare, U.F. Magaji, I.H. Abubakar, M. Yerima and R.S.U. Wasagu. (2018). Antifungal Potentials of *Acacia nilotica*, *Ziziphus jujube* Linn and *Lawsonia inermis*. *Asian Journal of Research in Biochemistry* 3 (1): 1-6.
- [41] T. Arif, J.D Bhosale, N. Kumar, T.K Mandal, R.S Bendre, G.S Lavekar and R. Dabur. (2009). Natural products antifungal agents derived from plants. *Journal of Asian Natural Products Research* 11 (7): 621-638.
- [42] M.M. Cowan. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 12 (4): 564–582.
- [43] A. Scalbert. (1991). Antimicrobial properties of tannins. *Phytochemistry* 30 (12): 3875–3883.
- [44] H.E. Brownlee, A.R. McEuen, J. Hedger and I.M. Scott. (1990). Anti-fungal effects of cocoa tannin on the witches' broom pathogen *Crinipellis pernicioso*. *Physiological and Molecular Plant Pathology* 36 (1): 39–48.
- [45] A. Bashir, I. Khan, S. Bashir, S. Azam and N. Ali. 2011. The antifungal, cytotoxic, antitermite, and insecticidal activities of *Ziziphus jujube*. *Pakistan Journal of Pharmaceutical Sciences* 24 (4): 489-493.
- [46] B. Khameneh, M. Iranshahy, V. Soheili and B.S. Fazly Bazzaz. (2019). Review on plant antimicrobials: a mechanistic viewpoint. *Antimicrobial Resistance and Infection Control* 8:118.
- [47] Z.B. Yi, Y. Yu, Y.Z. Liang and B. Zeng. (2007). Evaluation of the antimicrobial mode of berberine by LC/ ESI-MS combined with principal component analysis. *Journal of Pharmaceutical and Biomedical Analysis* 44 (1): 301–304.
- [48] D. Savoia. (2012). Plant-derived antimicrobial compounds: alternatives to antibiotics. *Future Microbiology* 7 (8): 979–990.
- [49] A.K. Agarwal, T. Xu, M.R Jacob, Q. Feng, M.C Lorenz, L.A Walker and A.M Clark. (2008). Role of heme in the antifungal activity of the azaxoaporphine alkaloid sampangine. *Eukaryotic Cell* 7(2):387-400.
- [50] L. Othman, A. Sleiman, R.M. Abdel-Massih. (2019). Antimicrobial Activity of Polyphenols and Alkaloids in Middle Eastern Plants. *Frontiers in Microbiology* 09: 911.